

Long-tailed Cormorant

Phalacrocorax africanus

738 plots from 1,964 records

Virgin Yellow-vented Bulbul squares **14**

Single YvB month **30**

Two YvB months **31**

Seasonality

	presence	bs	eggs
January	434	3	3
February	172	6	-
March	142	5	1
April	103	1	1
May	105	4	2
June	91	1	-
July	144	-	1
August	143	1	-
September	125	3	-
October	159	1	-
November	140	2	-
December	165	2	-
month 13	4		

Status


- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

Widespread and locally common, month gaps clearly show coverage problems. Rubondo gaps most surprising. Absence south of the Rufiji Delta is probably real. Although the Ruvouma is poorly worked this bird will not be common along this river.

Paucity of breeding records due to nesting low down in dense vegetation, needs effort to locate active nests.



with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
May 2007
From 762,609
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square