

# White-breasted Cormorant

*Phalacrocorax carbo*

334 plots from 846 records

Virgin Yellow-vented Bulbul squares **14**

Single YvB month **30**

Two YvB months **31**

Seasonality  
presence bs eggs


Month	presence	bs	eggs
January	214	2	7
February	82	2	1
March	57	2	2
April	34	3	-
May	32	2	-
June	42	1	-
July	49	1	-
August	60	4	-
September	59	4	-
October	58	2	-
November	52	2	-
December	68	1	1
month 13	2		

## Status

- Presence
- Breeding season
- Month with eggs

## Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900



Although widespread in low numbers there are only 4 documented breeding sites. That is if Lake Victoria can be considered a single site.

Formally common in Malawi now much reduced by fishermen and obviously rare along the northern shores of Lake Nyasi.

with support from  
THE LEVERHULME TRUST

**Tanzania Bird Atlas**  
**Preliminary Map**  
**May 2007**  
**From 762,609**  
**Database records**

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
within each square