

# Black River Duck

*Anas sparsa*

90 plots from 193 records

Virgin Yellow-vented Bulbul squares **20**

Single YvB month **37**

Two YvB months **34**

Seasonality  
presence bs eggs

Month	presence	bs	eggs
January	19	-	-
February	21	1	-
March	14	-	-
April	10	-	-
May	9	-	-
June	15	-	1
July	17	-	1
August	21	-	1
September	17	-	-
October	11	1	-
November	19	2	-
December	14	-	-

## Status

- Presence
- Breeding season
- Month with eggs

## Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

There are far fewer records here than there should be, This species will be largely resident on many (most !!) of the mountain streams in the Eastern Arc mts and Southern Highlands. It will also occur in the lower reaches of these streams.

It will also be on many of the streams in the far west. A record of a pair on a muddy stream near Biharamulo is shown by the blue circle.

with support from  
THE LEVERHULME TRUST

**Tanzania Bird Atlas**  
**Preliminary Map**  
**July 2005**  
**From 685,400**  
**Database records**

J	F	M	A
#	#	#	#
M	J	J	A
#	#	#	#
S	O	N	D
#	#	#	#

Monthly records  
within each square

