

Lappet-faced Vulture

Torgos tracheliotus

427 plots from 1,490 records

Virgin Yellow-vented Bulbul squares **14**

Single YvB month **30**

Two YvB months **31**

Seasonality
presence bs eggs


Month	presence	bs	eggs
January	184	3	2
February	139	4	2
March	111	5	4
April	95	4	5
May	72	5	4
June	108	4	3
July	132	5	2
August	140	11	2
September	92	2	-
October	126	4	-
November	88	2	-
December	127	1	-
month 13	2		

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900



with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
August 2007
From 767,217
Database records

J F M A
M J J A
S O N D
#

Monthly records
within each square

That this species is largely restricted to protected areas is both testament to the effectiveness of these sites and a clear indication that, rather sadly, it cannot exist outside the National Parks and Game Reserves and their stock of large mammals.

The lack of breeding data away from the northern protected areas is of concern. It should breed in these western, southern and eastern reserves.

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°