

Lammergeyer

Gypaetus barbatus

34 plots from 70 records

Virgin Yellow-vented Bulbul squares **33**

Single YvB month **32**

Two YvB months **30**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	6	-	-
February	4	-	-
March	6	-	-
April	2	-	-
May	-	-	-
June	4	-	-
July	9	-	-
August	11	-	-
September	6	-	-
October	9	-	-
November	6	-	-
December	7	-	-

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

This is a rare and highly restricted bird in Tanzania. These birds can be considered the southern limits of the population that extends southwards from Ethiopia through Kenya although it is now highly fragmented and gene flow in recent decades must be low.

It is only seen on a regular basis on mts Kilimanjaro and Meru where a small population of probably less than 50 birds is more or less resident. There are sufficient irregular sightings from the Gol Mts to suggest it is also possibly resident there and there are reports of a nest on the north side of Monduli Mt.

The single record from Mkomazi must be considered a wanderer from Kilimanjaro.

with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
January 2005
From 653,800
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

