

# Fulleborn's Black Boubou

*Laniarius fuelleborni*

131 plots plots from 289 records

Virgin Yellow-vented Bulbul squares **12**

Single YvB month **21**

Two YvB months **24**

Seasonality  
presence bs eggs

Month	presence	bs	eggs
January	33	5	-
February	42	8	1
March	25	1	-
April	30	-	-
May	28	1	-
June	35	-	-
July	45	-	-
August	25	1	-
September	34	2	-
October	46	2	-
November	43	6	-
December	38	9	-
month 13	4		

## Status

- Presence
- Breeding season
- Month with eggs

## 30m SRTM

- 850 - 900
- 900 - 950
- 950 - 1,000
- 1,000 - 1,050
- 1,050 - 1,100
- 1,100 - 1,200
- 1,200 - 1,300
- 1,300 - 1,400
- 1,400 - 1,500
- 1,500 - 1,600
- 1,600 - 1,700
- 1,700 - 1,800
- 1,800 - 1,900
- 1,900 - 2,000
- 2,000 - 2,100
- 2,100 - 2,200
- 2,200 - 2,300
- 2,300 - 2,400
- 2,400 - 2,500

Restricted to montane forest in most of the Eastern Arc mts. The race *L.f. usambaricu* is known from the West Usambara, Nguru, Ukaguru and Uluguru mts but not the East Usambara. The nominate race occurs in the Udzungwa and further south.

Tanzania Bird Atlas  
Preliminary Map  
September 2009  
From 862,509  
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
within each square

