

Olive-bellied Sunbird

Nectarinia chloropygia

29 plots from 63 records

Virgin Yellow-vented Bulbul squares **12**

Single YvB month **21**

Two YvB months **24**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	3	-	-
February	5	-	-
March	2	-	-
April	3	-	-
May	3	-	-
June	2	-	-
July	10	-	-
August	8	-	-
September	6	-	-
October	5	-	-
November	12	1	1
December	2	-	-

Status

- Presence
- Breeding season
- Month with eggs

30m SRTM

- 0 - 100
- 100 - 200
- 200 - 300
- 300 - 400
- 400 - 450
- 450 - 500
- 500 - 550
- 550 - 600
- 600 - 650
- 650 - 700
- 700 - 750
- 750 - 800
- 800 - 850
- 850 - 900
- 900 - 950
- 950 - 1,000
- 1,000 - 1,050
- 1,050 - 1,100
- 1,100 - 1,200
- 1,200 - 1,300
- 1,300 - 1,400
- 1,400 - 1,500
- 1,500 - 1,600
- 1,600 - 1,700
- 1,700 - 1,800
- 1,800 - 1,900
- 1,900 - 2,000
- 2,000 - 2,100
- 2,100 - 2,200
- 2,200 - 2,300
- 2,300 - 2,400
- 2,400 - 2,500

A West African species restricted in Tanzania to the far west. It should occur in forest edge and riverine throughout this range but no further east than 32 degrees.

The paucity of data reflects the absence of resident birders throughout these border areas.

One would have expected more records from Rubondo NP and Mahale NP where it is surely overlooked.

Tanzania Bird Atlas
Preliminary Map
August 2009
From 861,488
Database records

J F M A
M J J A
S O N D
#

Monthly records
within each square

