

Red-chested Sunbird

Nectarinia erythroceria

134 plots from 260 records

Virgin Yellow-vented Bulbul squares **12**

Single YvB month **21**

Two YvB months **24**

Seasonality

	presence	bs	eggs
January	27	1	-
February	22	3	1
March	21	-	2
April	13	-	-
May	18	-	1
June	17	-	-
July	29	-	-
August	15	1	-
September	16	1	-
October	27	-	-
November	15	1	-
December	24	3	-
month 13	2		

Restricted to waterside habitats including riverine within its range. The gap between Mahale NP and Kigoma and further north is most probably an artifact of poor coverage.

It is particularly common along the shore of Lake Victoria, even in drier habitats along the eastern shoreline.


Britton (1980) also includes Kahama (3203D) but this is well south of the range mapped here.

Status

- Presence
- Breeding season
- Month with eggs

30m SRTM

- 0 - 100
- 100 - 200
- 200 - 300
- 300 - 400
- 400 - 450
- 450 - 500
- 500 - 550
- 550 - 600
- 600 - 650
- 650 - 700
- 700 - 750
- 750 - 800
- 800 - 850
- 850 - 900
- 900 - 950
- 950 - 1,000
- 1,000 - 1,050
- 1,050 - 1,100
- 1,100 - 1,200
- 1,200 - 1,300
- 1,300 - 1,400
- 1,400 - 1,500
- 1,500 - 1,600
- 1,600 - 1,700
- 1,700 - 1,800
- 1,800 - 1,900
- 1,900 - 2,000
- 2,000 - 2,100
- 2,100 - 2,200
- 2,200 - 2,300
- 2,300 - 2,400
- 2,400 - 2,500



Tanzania Bird Atlas
Preliminary Map
August 2009
From 861,488
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records within each square