

# Augur Buzzard

*Buteo augur*

859 plots from 2,924 records

Locally common, easily identified and well adapted to human environments where it does rather well on rodents inhabiting agricultural land. It is likely to be more widespread within the Miombo woodlands than indicated here. Its absence from the coastal zone is well known.

Virgin Yellow-vented Bulbul squares **33**

Single YvB month **34**

Two YvB months **30**

Seasonality  
presence bs eggs


Month	presence	bs	eggs
January	300	4	1
February	254	1	-
March	221	8	2
April	209	6	1
May	164	10	1
June	189	11	4
July	275	7	7
August	267	8	2
September	219	8	-
October	247	6	1
November	219	3	2
December	261	7	1

## Status

- Presence
- Breeding season
- Month with eggs

## Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900



with support from  
THE LEVERHULME TRUST

Tanzania Bird Atlas  
Preliminary Map  
December 2004  
From 651,000  
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
within each square

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°

-1°  
-2°  
-3°  
-4°  
-5°  
-6°  
-7°  
-8°  
-9°  
-10°  
-11°