

Steppe Buzzard

Buteo buteo vulpinus

437 plots from 995 records

Virgin Yellow-vented Bulbul squares **33**

Single YvB month **34**

Two YvB months **30**

Seasonality presence bs eggs

Month	presence	bs	eggs
January	155	-	-
February	180	-	-
March	108	-	-
April	34	-	-
May	5	-	-
June	6	-	-
July	5	-	-
August	6	-	-
September	24	-	-
October	122	-	-
November	186	-	-
December	162	-	-

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900


A locally common bird on passage where it can occur virtually anywhere. It is a regular winter visitor to highland areas in southern Tanzania but less common in the north.

with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
December 2004
From 651,000
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square



29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°