

Lizard Buzzard

Kaupifalco monogrammicus

742 plots from 1,425 records

Virgin Yellow-vented Bulbul squares **13**

Single YvB month **25**

Two YvB months **28**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	94	-	-
February	112	-	1
March	85	1	-
April	78	-	-
May	96	1	-
June	121	-	-
July	183	2	-
August	147	5	-
September	144	2	-
October	103	3	-
November	105	1	1
December	126	2	1
month 13	11		

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)


- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

Mostly resident in higher rainfall areas but evidently wanders more widely after the dry season, post breeding dispersal !! At the level of the Atlas square this species utilises virtually all non forested habitats except the driest.

Tanzania Bird Atlas
Preliminary Map
April 2008
From 803,703
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square



29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°