

# Gabar Goshawk

*Melierax gabar*

614 plots from 1,261 records

Virgin Yellow-vented Bulbul squares **33**

Single YvB month **34**

Two YvB months **30**

Seasonality  
presence bs eggs

Month	presence	bs	eggs
January	127	1	-
February	105	4	-
March	110	3	-
April	93	1	-
May	95	-	-
June	110	3	-
July	116	-	-
August	133	1	-
September	100	-	-
October	77	-	-
November	92	-	-
December	87	2	-

## Status

- Presence
- Breeding season
- Month with eggs

## Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900


Widespread and locally common in drier woodland, less so in the coastal lowlands and quite rare in the west where it probably competes more with the Shrika.

with support from  
THE LEVERHULME TRUST

Tanzania Bird Atlas  
Preliminary Map  
December 2004  
From 651,000  
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
within each square



29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°

-1°  
-2°  
-3°  
-4°  
-5°  
-6°  
-7°  
-8°  
-9°  
-10°  
-11°