

Black-shouldered Kite

Elanus caeruleus

1,232 plots from 3,508 records

Widespread and locally common but we know almost nothing about movements other than obvious invasions during times of prey abundance. Eggs produced in most months is also unusual, no obvious seasonality !!

Virgin Yellow-vented Bulbul squares **14**

Single YvB month **30**

Two YvB months **31**


Seasonality	presence		bs	eggs
	bs	eggs		
January	404	2	3	
February	326	2	-	
March	244	8	3	
April	215	2	4	
May	196	5	3	
June	249	2	3	
July	357	4	2	
August	315	5	2	
September	243	3	1	
October	281	4	-	
November	299	4	1	
December	313	1	1	
month 13	1			

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900



with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
August 2007
From 767,217
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°

-1°
-2°
-3°
-4°
-5°
-6°
-7°
-8°
-9°
-10°
-11°