

# Hartlaub's Bustard

*Eupodotis hartlaubi*

82 plots from 150 records

Virgin Yellow-vented Bulbul squares **18**

Single YvB month **37**

Two YvB months **34**

### Seasonality

	presence	bs	eggs
January	15	3	-
February	15	1	-
March	5	3	-
April	13	4	-
May	6	2	-
June	8	-	-
July	6	1	-
August	17	-	-
September	11	-	-
October	6	-	-
November	10	1	-
December	19	3	-

### Status

- Presence
- Breeding season
- Month with eggs

### Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

An uncommon species largely restricted to drier highland areas.

The 6 records at the west and southern periphery of the range are all old ones from before 1981.

Unless seen well this species can be confused with the much more common Black-bellied and as such may be under recorded.

with support from  
THE LEVERHULME TRUST

**Tanzania Bird Atlas**  
**Preliminary Map**  
**September 2005**  
**From 690,800**  
**Database records**

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records within each square

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°

