

Black-bellied Bustard

Eupodotis melanogaster

465 plots from 908 records

Virgin Yellow-vented Bulbul squares **18**

Single YvB month **37**

Two YvB months **34**

Seasonality

	presence	bs	eggs
January	102	12	1
February	103	7	-
March	79	1	2
April	64	5	-
May	38	2	3
June	54	1	1
July	67	1	-
August	57	-	-
September	59	-	-
October	69	1	-
November	82	5	-
December	87	4	-
month 13	1		

Status


- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

Widespread and often locally common. In general terms the paucity of records in the west and SE reflects poor coverage in these areas.

Regular movements occur, probably in response to rainfall but the data as not yet been analysed.



with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
September 2005
From 690,800
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°

-1°
-2°
-3°
-4°
-5°
-6°
-7°
-8°
-9°
-10°
-11°