

Kori Bustard

Ardeotis kori

283 plots from 1,127 records

Virgin Yellow-vented Bulbul squares **18**

Single YvB month **37**

Two YvB months **34**

Seasonality


	presence	bs	eggs
January	156	13	1
February	96	11	1
March	84	14	4
April	72	12	3
May	63	1	-
June	72	-	-
July	87	1	-
August	91	-	-
September	58	-	-
October	83	1	-
November	68	4	1
December	117	12	-
month 13	1		

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900



Largely restricted to the dry northern grasslands where it is common in protected areas but far less so elsewhere.

There is talk of birds being seen in Ruaha NP and the records from Mtera indicate there could be a small population in this general area.

The odd records from Rubondo NP and other areas west of the main range indicate seasonal wandering. Its absence from the Wembere Steppe is surprising.

This population has been repeatedly plundered for the international zoo trade and in some areas it is now far less common than even two decades ago. It is also hunted indiscriminately, witness several birds shot by the northern zone anti-poaching unit in the mid 90s ostensibly to feed visiting government dignitaries .

with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
September 2005
From 690,800
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square