

It is still far from clear if we have an influx of this bird at any time and, if we do, where they might originate from. It appears largely resident in all well watched sites in the north but far less so in the south.

Curious that there are, as yet, no breeding records from the Selous and Kilombero river systems.

Kittlitz's Plover

Charadrius pecuarius

421 plots from 1,275 records

Virgin Yellow-vented Bulbul squares **13**

Single YvB month **25**

Two YvB months **28**


	Seasonality		
	presence	bs	eggs
January	186	-	4
February	109	1	4
March	92	5	7
April	77	2	4
May	61	5	4
June	62	5	18
July	100	8	18
August	99	5	3
September	87	4	11
October	83	4	5
November	71	2	5
December	116	2	4
month 13	2		

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- VALUE**
- 1 - 300
 - 300 - 500
 - 500 - 700
 - 700 - 900
 - 900 - 1,100
 - 1,100 - 1,300
 - 1,300 - 1,500
 - 1,500 - 1,700
 - 1,700 - 1,900
 - >1,900



Tanzania Bird Atlas
Preliminary Map
June 2008
From 809,297
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square