

Wattled Plover

Vanellus senegallus

408 plots from 828 records

Virgin Yellow-vented Bulbul squares **13**

Single YvB month **25**

Two YvB months **28**

Seasonality

	presence	bs	eggs
January	115	2	4
February	73	2	2
March	61	-	1
April	32	2	-
May	39	1	-
June	54	3	1
July	86	2	-
August	68	1	4
September	60	4	6
October	60	4	3
November	62	4	5
December	59	3	3
month 13	2		

Species

- Presence
- Breeding season
- Eggs in nest

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

There is a suggestion that this bird is expanding its range eastwards into north-central & central areas and into the Selous GR. 30 years ago the few localities for Tanzania (6 mentioned) were discussed in the first two issues of Scopus.

It is now known from 96 (27%) Atlas squares.


The 1% threshold for this population is 1,000 birds, the only likely "site" in Tanzania that qualifies is the Moyowosi-Malagarasi Ramsar site.

The draft Wader Atlas maps sites of 2-50, 51-100 & 100-216 birds with two of the smaller and a single >100 site in the Kilombero Valley. Only 82 were counted nationwide in January 2005.

Tanzania Bird Atlas
Preliminary Map
June 2008
From 809,297
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square



-1°
-2°
-3°
-4°
-5°
-6°
-7°
-8°
-9°
-10°
-11°

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°