

Spur-winged Plover

Vanellus spinosus

291 plots from 605 records

The 1% threshold for this species is 4,000 birds, there are no such sites in Tanzania.

The draft Wader Atlas maps sites for concentrations of 99-500, 501-1,000 and 1001-2753 birds. No sites are mapped for Tanzania but concentrations exceeding 100 birds can now be expected at favoured sites such as Nyumba ya Mungu.

Virgin Yellow-vented Bulbul squares **13**

Single YvB month **25**

Two YvB months **28**

Seasonality
presence bs eggs


Month	presence	bs	eggs
January	121	1	2
February	50	1	2
March	40	-	-
April	29	1	-
May	37	-	1
June	37	-	-
July	51	-	2
August	53	4	2
September	44	2	4
October	42	1	2
November	32	-	-
December	40	2	1
month 13	1		

Species

- Presence
- Breeding season
- Eggs in nest

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900



Tanzania Bird Atlas
Preliminary Map
June 2008
From 809,297
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

The southern range expansion of this species continues unabated. There are now odd records from Zambia and Malawi and many more sites within Tanzania are being utilised by this bird. Britton (1980) states "There is no breeding record for Tanzania where all or most records are likely to refer to wanderers, south to Morogoro and Rukwa". Baker (1994) listed a further 10 sites including the first two breeding records. It now occurs in 82 (23%) squares and can be expected to continue increasing.