

Eurasian Curlew

Numenius arquata

125 plots from 251 records

Virgin Yellow-vented Bulbul squares **13**

Single YvB month **25**

Two YvB months **28**

Seasonality
presence bs eggs

January	77	-	-
February	26	-	-
March	21	-	-
April	12	-	-
May	5	-	-
June	3	-	-
July	14	-	-
August	16	-	-
September	16	-	-
October	15	-	-
November	20	-	-
December	24	-	-
month 13	2	-	-

status


● Presence

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

A uncommon bird at these latitudes with small but regular over wintering populations along the coast, esp south to Mafia, and in the Northern Rift centred on the Serengeti and Ngorongoro. Only scattered records elsewhere. Inland and coastal populations may have different origins.

Only 28 birds were counted by 4 teams in 1995 and 419 by 8 teams in 2005. It is unlikely that Tanzania holds many more than 1,000 birds.



Tanzania Bird Atlas
Preliminary Map
June 2008
From 809,297
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square