

Whimbrel

Numenius phaeopus

151 plots from 605 records

Virgin Yellow-vented Bulbul squares **13**

Single YvB month **25**

Two YvB months **28**

Seasonality	presence		bs	eggs
	presence	bs		
January	114	-	-	-
February	50	-	-	-
March	54	-	-	-
April	38	-	-	-
May	19	-	-	-
June	34	-	-	-
July	35	-	-	-
August	47	-	-	-
September	55	-	-	-
October	52	-	-	-
November	42	-	-	-
December	65	-	-	-

status

● Presence

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

This is a common bird along our coast as far south as Mafia but distinctly less so further south and is rare inland. Only 3,383 were counted in January 2005 and the more detailed Danish survey in 1989 counted 3,872 along 127km of coast.

The draft Wader Atlas maps sites of 231-1,000, 1001-3000 and 3,001-6,000. Only such large "sites" as Pemba, Dar coast, Mafia, Rufiji Delta & Mnazi Bay qualify at present.

**Tanzania Bird Atlas
Preliminary Map
June 2008
From 809,297
Database records**

J F M A
M J J A
S O N D
#

Monthly records
within each square

