

Greenshank

Tringa nebularia

640 plots from 1,973 records

There are no 1% sites (10,000 birds) for this population throughout its vast range. The draft Wader Atlas maps sites of 300-600, 601-1500 & 1501-3200. There are none of the latter in Africa, only 8 of the mid range and 8 smaller sites. This suggests serious under counting and / or an overestimate of the population size.

Only 2,407 were counted by 16 teams in Jan 1995 and 2,626 by 24 teams in Jan 2005. Baker (1996) suggested a max mid winter population of 10,000. It could be twice this.

This species prefers the shores of the larger shallow waterbodies and the coast with smaller numbers along the larger rivers.

Virgin Yellow-vented Bulbul squares **13**

Single YvB month **25**

Two YvB months **28**


	Seasonality		
	presence	bs	eggs
January	430	-	-
February	197	-	-
March	174	-	-
April	91	-	-
May	48	-	-
June	52	-	-
July	102	-	-
August	161	-	-
September	178	-	-
October	189	-	-
November	149	-	-
December	199	-	-
month 13	3		

status

● Presence

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900



Tanzania Bird Atlas
Preliminary Map
June 2008
From 809,297
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square