

This is not a congregatory species and prefers muddy pools and the smaller fast flowing rivers and streams. The draft Wader Atlas maps concentrations of 27-100, 101-300 and 301-818. There are only 8 of the smaller sites in Africa for this eastern population and a single mid range site. The population estimate is 100,000 to 1M. If this site system followed the IBA approach then many of our NPs and GRs should be mapped.

Green Sandpiper

Tringa ochropus

408 plots from 986 records

- Virgin Yellow-vented Bulbul squares **13**
- Single YvB month **25**
- Two YvB months **28**


	Seasonality		
	presence	bs	eggs
January	263	-	-
February	143	-	-
March	81	-	-
April	13	-	-
May	8	-	-
June	4	-	-
July	10	-	-
August	42	-	-
September	64	-	-
October	112	-	-
November	100	-	-
December	145	-	-
month 13	1	-	-

status

● Presence

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900



Tanzania Bird Atlas
Preliminary Map
June 2008
From 809,297
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square