

Terek Sandpiper

Xenus cinereus

112 plots from 353 records

Virgin Yellow-vented Bulbul squares **13**

Single YvB month **25**

Two YvB months **28**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	87	-	-
February	33	-	-
March	36	-	-
April	26	-	-
May	11	-	-
June	15	-	-
July	5	-	-
August	11	-	-
September	26	-	-
October	34	-	-
November	31	-	-
December	38	-	-

status

● Presence

Elevation (m)

1 - 300

300 - 500

500 - 700

700 - 900

900 - 1,100

1,100 - 1,300

1,300 - 1,500

1,500 - 1,700

1,700 - 1,900

>1,900

Rare inland but locally common along the coast, more so north of the Rufiji than to the south.

Bregnballe (1989) counted 2,629 birds along 127km of coastline. Only 178 were counted by 2 teams in January 1995. 1,605 were counted by 7 teams in January 2005. No 1% (10,000 birds) sites are known throughout the range of the species.. The draft Wader Atlas maps 425-500, 501-1,500 and 1501-2895 sites. There are several sites mapped for Tanzania.

Tanzania Bird Atlas
Preliminary Map
June 2008
From 809,297
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

