

Curlew Sandpiper

Calidris ferruginea

326 plots from 951 records

Virgin Yellow-vented Bulbul squares **13**

Single YvB month **25**

Two YvB months **27**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	182	-	-
February	87	-	-
March	79	-	-
April	86	-	-
May	42	-	-
June	25	-	-
July	35	-	-
August	93	-	-
September	83	-	-
October	92	-	-
November	68	-	-
December	79	-	-

status

● Presence

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

Wdespread inland but large numbers occur only along the coast. 2004 & 8636 on Pemba & Zanzibar in 1998. 16,158 by 10 teams in Jan 95 included 7,000 in the northern Rift and 6,000 along the Dar es Salaam coast. Only 6,515 in January 2005 but the Rufiji Delta was missed and this held 16,000 birds in 2000. Bregnballe (1989) counted almost 15,000 birds along the coast between Dar es Salaam and Kilwa in January 1988 at densities of 10.4 - 647.3 per km.

The 1% threshold is 4,000 birds and the draft Wader Atlas maps concentrations from 3205 to a staggering 505,000 birds on the Bijagos Archipelago in Guinea-Bissau.

Tanzania Bird Atlas
Preliminary Map
June 2008
From 811,235
Database records

J F M A
M J J A
S O N D
#

Monthly records
within each square

