

Whiskered Tern

Chlidonias hybridus

293 plots from 679 records

Virgin Yellow-vented Bulbul squares **14**

Single YvB month **30**


Two YvB months **31**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	126	20	1
February	49	24	-
March	53	15	2
April	34	16	1
May	21	12	1
June	33	5	1
July	38	6	-
August	36	4	-
September	33	3	-
October	40	4	-
November	31	8	-
December	49	12	-

Status

- Presence
- Breeding season
- Month with eggs



Britton (1980) considers all birds to be the race *delalandii* with no evidence that Palearctic birds reach this far south.

There are only four breeding sites shown here simply because it is often difficult to actually prove egg laying due to access difficulties.

The majority of breeding season records relate to birds being seen in breeding plumage in suitable habitat. However they are known to undertake considerable movements so one must take care with such observations. Perhaps the best indication of breeding is the regular movement of birds to and from a given location, especially when they can be seen carrying food.

There will be many more breeding sites than shown here and locating these is something of a priority. Local canoes are ideal on most waters and it is never too cold not to swim !

with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
January 2007
From 751,930
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°