

# Grey Go-away Bird

*Corythaixoides concolor*

85 plots from 131 records

Virgin Yellow-vented Bulbul squares **20**

Single YvB month **37**

Two YvB months **34**

	Seasonality		
	presence	bs	eggs
January	10	-	-
February	9	-	-
March	10	-	-
April	7	-	-
May	14	-	-
June	15	-	-
July	13	-	-
August	11	-	-
September	13	-	-
October	12	-	-
November	9	-	-
December	6	1	-
month 13	1		

### Status

- Presence
- Breeding season
- Month with eggs

### Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

This southern African species reaches its northern limits in Tanzania.

Its restricted distribution is somewhat surprising as there is no obvious competitor throughout the coastal lowlands. It does not occur in Miombo Woodland and this limits its range towards the higher ground in the southern Selous and beyond.

The few records in the south and along the valley of the Ruvuma River are indicative of poor observer coverage in this general area but it is not at all a common bird here.

There are probably links with a population in northern Mocambique but this area is also under recorded.

A recent record from just south of Dar es Salaam as not yet been entered in the database.

with support from  
THE LEVERHULME TRUST

**Tanzania Bird Atlas**  
**Preliminary Map**  
**August 2005**  
**From 688,400**  
**Database records**

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
within each square

