

Bare-faced Go-away Bird

Corythaixoides personata

737 plots from 1,499 records

Virgin Yellow-vented Bulbul squares **20**

Single YvB month **37**

Two YvB months **34**

Seasonality


	presence	bs	eggs
January	138	2	-
February	124	1	1
March	104	1	-
April	100	2	-
May	77	1	1
June	144	3	-
July	158	-	-
August	145	-	2
September	130	-	-
October	137	1	-
November	117	-	-
December	110	-	-

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900



with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
August 2005
From 688,400
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

Locally common throughout the western Miombo woodlands this species shows considerable penetration into the drier range of White-bellied Go-away Bird.

Its apparent absence in the wetter environment of the far west awaits confirmation from this underworked area.

Despite being highly visible there are very few breeding records for any of the go-away birds.

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°