

Livingstone's Turaco

Turaco livingstonii

241 plots from 512 records

Virgin Yellow-vented Bulbul squares **33**

Single YvB month **31**

Two YvB months **31**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	47	-	-
February	50	5	-
March	40	1	-
April	24	-	-
May	26	-	-
June	23	-	-
July	38	-	-
August	33	1	-
September	51	1	-
October	52	-	1
November	63	1	-
December	53	2	-

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

A bird of the coastal forests south of the Wami River and the montane forests of the Eastern Arc mts and those around the north-western end of Lake Nyasa and Mbeya whence it extends south into Malawi and Mocambique.

The isolated record from 3705A is from the Nguu Mts and this species will occur in the squares immediately to the east, south and south-east. This montane forest represents the northern limits of this species.


There is a curious specimen record in the literature from well within the range of Schalow's Turaco near Biharamulo in the north-west, shown here as a red circle. The origins and affinities of this specimen remain a mystery.

with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
March 2005
From 667,000
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square



29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°

-1° -2° -3° -4° -5° -6° -7° -8° -9° -10° -11°