

# Black-billed Turaco

*Tauraco schuetti*

12 plots from 18 records

Virgin Yellow-vented Bulbul squares **33**

Single YvB month **31**

Two YvB months **31**

Seasonality  
presence bs eggs

Month	presence	bs	eggs
January	-	-	-
February	-	-	-
March	-	-	-
April	-	-	-
May	-	-	-
June	1	-	-
July	4	-	-
August	4	-	-
September	3	-	-
October	3	-	-
November	2	1	-
December	-	-	-

## Status

- Presence
- Breeding season
- Month with eggs

## Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

In Tanzania this species is only known from Minziro FR on the border with Uganda.

The two records from Mahale NP are both recent and from the same observer. They require confirmation before acceptance.

with support from  
THE LEVERHULME TRUST

**Tanzania Bird Atlas**  
**Preliminary Map**  
**March 2005**  
**From 667,000**  
**Database records**

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
within each square

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°

