

Black Cuckoo

Cuculus clamosus

315 plots from 514 records

Virgin Yellow-vented Bulbul squares **13**

Single YvB month **25**

Two YvB months **28**

Seasonality

	presence	bs	eggs
January	105	8	-
February	68	11	-
March	58	4	-
April	33	3	-
May	22	2	-
June	6	-	-
July	4	-	-
August	3	-	-
September	7	-	-
October	34	2	-
November	53	7	-
December	74	8	-
month 13	2		

Status


- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

Does this bird leave Tanzania or is it just too difficult to locate when not calling ?

Grateful for a scan of the latest text on this species from both of the recent cuckoo books.



with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
May 2008
From 807,213
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°