

Red-chested Cuckoo

Cuculus solitarius

609 plots from 1,596 records

Virgin Yellow-vented Bulbul squares **18**

Single YvB month **37**

Two YvB months **34**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	272	8	1
February	198	5	1
March	130	3	-
April	115	1	-
May	69	5	1
June	42	1	-
July	29	1	-
August	28	-	-
September	46	-	-
October	130	2	6
December	291	5	-
month 13	3		

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900


Difficult to locate when not calling and too easy when it is !! Therefore the bias towards rainy season records may not accurately reflect the true distributional seasonality although resident calling birds will indicate breeding season.

with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
September 2005
From 690,800
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square



29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°