

Mottle-throated Spinetail

Telacanthura ussheri

213 plots from 424 records

Virgin Yellow-vented Bulbul squares **34**

Single YvB month **38**

Two YvB months **35**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	41	1	-
February	41	3	-
March	31	1	-
April	22	-	-
May	17	1	-
June	38	1	-
July	38	-	-
August	35	-	-
September	33	-	-
October	48	1	-
November	23	1	-
December	46	1	-

Status

- Presence
- Breeding season
- Month with eggs

Protected Areas

- National Park
- Game Reserve
- Conservation Area

Elevation (m)

- 1 - 500
- 500 - 800
- 800 - 1,100
- 1,100 - 1,400
- 1,400 - 1,700
- 1,700 - 1,900

A widespread but local species on the central plateau where it is closely associated with Baobabs in which it breeds. It is also local along the coast where it is loosely associated with coastal forest. The few records from the north-west are of the West African race *T. u. sharpei*, a forest bird known in Tanzania only from Minziro FR.

Tanzania Bird Atlas
Preliminary Map
October 2004
From 641,500
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

