

Blue-breasted Kingfisher

Halcyon malimbica

38 plots from 69 records

Virgin Yellow-vented Bulbul squares **12**

Single YvB month **21**

Two YvB months **24**

Seasonality

presence bs eggs

Month	presence	bs	eggs
January	6	-	-
February	4	-	-
March	3	-	-
April	1	-	-
May	2	-	-
June	2	-	-
July	15	-	-
August	8	-	-
September	3	-	-
October	7	-	-
November	12	1	-
December	5	-	-

Status

- Presence
- Breeding season
- Month with eggs

30m SRTM

- 0 - 100
- 100 - 200
- 200 - 300
- 300 - 400
- 400 - 450
- 450 - 500
- 500 - 550
- 550 - 600
- 600 - 650
- 650 - 700
- 700 - 750
- 750 - 800
- 800 - 850
- 850 - 900
- 900 - 950
- 950 - 1,000
- 1,000 - 1,050
- 1,050 - 1,100
- 1,100 - 1,200
- 1,200 - 1,300
- 1,300 - 1,400
- 1,400 - 1,500
- 1,500 - 1,600
- 1,600 - 1,700
- 1,700 - 1,800
- 1,800 - 1,900
- 1,900 - 2,000
- 2,000 - 2,100
- 2,100 - 2,200
- 2,200 - 2,300
- 2,300 - 2,400
- 2,400 - 2,500

The apparent gaps in the distribution of this species are clearly related to the paucity of records from western Tanzania. The month gaps in Mahale NP are curious as this is the one locality with resident observers and the gaps do not suggest seasonality / movements. It was in 3005C when we visited in August and November. It surely occurs along all well wooded streams and forest edge habitat throughout this range.

Tanzania Bird Atlas
Preliminary Map
August 2009
From 861,488
Database records

J F M A
M J J A
S O N D
#

Monthly records within each square

