

White-throated Bee-eater

Merops albicollis

264 plots from 534 records

Virgin Yellow-vented Bulbul squares **33**

Single YvB month **32**

Two YvB months **30**

Seasonality presence bs eggs

Month	presence	bs	eggs
January	101	1	-
February	81	-	-
March	82	-	-
April	35	1	-
May	9	1	-
June	6	-	-
July	14	-	-
August	14	-	-
September	16	-	-
October	31	-	-
November	51	-	-
December	87	1	-
undated	1		

Status

- ! Presence
- (Breeding season
- (Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

There is a small (numbers unknown) breeding population in the Natron Rift extending northwards to Lake Magadi.

The breeding season records from the Selous and near Bukoba probably refer to northern breeders displaying before migration. These 2 records are omitted from the seasonality table.

We need many more breeding season records of this species from the Natron Rift to firmly establish the status of this isolated population.

with support from THE LEVERHULME TRUST

Tanzania Bird Atlas Preliminary Map March 2006 From 722,000 Database records

J F M A
M J J A
S O N D
#

Monthly records within each square

