

Red-billed Hornbill

Tockus erythrorhynchus

238 plots from 795 records

Virgin Yellow-vented Bulbul squares **12**

Single YvB month **25**

Two YvB months **25**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	88	3	1
February	65	3	-
March	51	2	-
April	59	1	-
May	43	1	-
June	51	-	-
July	83	-	1
August	85	-	-
September	56	-	-
October	71	-	-
November	45	-	1
December	81	3	1

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

To accurately define the range of this species, especially with reference to Ruaha Hornbill we need GIS data from all western and SW border squares. The suspicion is that this species only occurs east of the Ruft Valley wall, east of the Mbulu Highlands and north of the Kondoa-Kibaya Ridge. There are a few records to clarify on the edge of this range that could be good for either species.

3504A Dongobesh-Haydom Dave Peterson
Katesh Paul Oliver

3504C Katesh Marc Baker

3605C Zoissa Bruce Hills

3706C Kimamba John Beesley

GPS records from ALL squares marked with GREEN circles PLEASE.

Tanzania Bird Atlas
Preliminary Map
January 2009
From 828,055
Database records

J F M A
M J J A
S O N D
#

Monthly records
within each square

