

Black-throated Barbet

Lybius melanocephalus

168 plots from 250 records

Virgin Yellow-vented Bulbul squares **33**

Single YvB month **32**

Two YvB months **30**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	16	1	-
February	15	1	1
March	12	1	-
April	16	-	-
May	18	2	-
June	17	1	-
July	30	1	-
August	27	-	-
September	18	1	-
October	17	-	1
November	17	1	1
December	30	5	-

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

These few records represent the endemic race *L. m. flavibuccalis* around the Wembere Steppe.

THIS MAP IS FOR GENERAL INTEREST ONLY AS IT CONTAINS SEVERAL ERRORS.

RECORDS LIKELY TO BE REJECTED ARE SHOWN IN RED CIRCLES. OTHER RECORDS REQUIRING VERIFICATION ARE SHOWN IN BLUE CIRCLES.

with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
December 2004
From 653,000
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°

