

A widespread bird of forest interior, forest edge, riverine, wet thicket, (dry thicket in Itigi) and rich Miombo. Difficult to locate when not calling but easily caught in mist nets.

Britton (1980) allows 3 races in Tanzania. Named localities as marked by symbols.

*S. c. suahelicus* in the coastal lowlands (red circles)  
*S. c. medianus* is wholly montane from the Usambara to Kilimanjaro and the Crater Highlands (blue squares)  
*S. c. albigularis* in western and central areas (green diamonds).

# African Broadbill

*Smithornis capensis*

257 plots from 633 records

Virgin Yellow-vented Bulbul squares **14**


Single YvB month **30**

Two YvB months **31**

	Seasonality		
	presence	bs	eggs
January	39	17	6
February	44	11	3
March	49	11	2
April	13	1	1
May	34	4	2
June	36	-	-
July	63	2	-
August	45	-	-
September	57	5	1
October	51	4	3
November	77	12	7
December	24	3	1
month 13	4	1	

## Status

- Presence
- Breeding season
- Month with eggs



with support from  
THE LEVERHULME TRUST

Tanzania Bird Atlas  
Preliminary Map  
January 2007  
From 751,930  
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
within each square