

# Black Roughwing

*Psalidoprocne holomelas*

620 plots from 1,794 records

Virgin Yellow-vented Bulbul squares **20**

Single YvB month **37**

Two YvB months **34**

Seasonality  
presence bs eggs

January	71	3	-
February	60	1	-
March	40	-	-
April	23	-	-
May	24	-	-
June	18	-	-
July	24	-	-
August	25	-	-
September	30	-	-
October	39	-	-
November	73	1	1
December	78	-	-
month 13	1		

Britton (1980) was adopted as the nomenclature for the Atlas in the early 1980s. This work did not split the Eastern Roughwing *Psalidoprocne orientalis* from the *P. pristoptera* / *holomelas* / *massaica* group.

Since the late 90s we have separated records of *orientalis* and these are shown on this map.

However, we still do not fully understand the species / subspecies limits of this group as only adult males are separable in the field.

● Eastern Roughwing

● Black Roughwing

● Presence

● Breeding season

● Month with eggs

with support from  
THE LEVERHULME TRUST

**Tanzania Bird Atlas**  
**Preliminary Map**  
**July 2005**  
**From 685,400**  
**Database records**

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
within each square

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°

