

Black Cuckoo Shrike

Campephaga flava

748 plots from 1,644 records

One of our most widespread and familiar birds but only in 3 squares do we have records for every month of the year. An analysis of movements is long overdue, any volunteers ? 1,644 records should tell us something.

Considered an Intra-African migrant and breeding visitor in Zambia (91% of squares). 73% coverage in Malawi "wanders widely Apr to Oct". No breeding records Uganda nearly all records Feb-Sept.

ITS COMPLICATED

Virgin Yellow-vented Bulbul squares **12**

Single YvB month **21**


Two YvB months **24**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	150	8	1
February	144	7	1
March	139	2	1
April	82	2	1
May	91	1	2
June	141	1	-
July	167	2	1
August	144	-	-
September	136	1	-
October	128	3	-
November	136	-	-
December	144	4	2
month 13	2		

- Status**
- Presence
 - Breeding season
 - Month with eggs

- 30m SRTM**
- 0 - 100
 - 100 - 200
 - 200 - 300
 - 300 - 400
 - 400 - 450
 - 450 - 500
 - 500 - 550
 - 550 - 600
 - 600 - 650
 - 650 - 700
 - 700 - 750
 - 750 - 800
 - 800 - 850
 - 850 - 900
 - 900 - 950
 - 950 - 1,000
 - 1,000 - 1,050
 - 1,050 - 1,100
 - 1,100 - 1,200
 - 1,200 - 1,300
 - 1,300 - 1,400
 - 1,400 - 1,500
 - 1,500 - 1,600
 - 1,600 - 1,700
 - 1,700 - 1,800
 - 1,800 - 1,900
 - 1,900 - 2,000
 - 2,000 - 2,100
 - 2,100 - 2,200
 - 2,200 - 2,300
 - 2,300 - 2,400
 - 2,400 - 2,500



Tanzania Bird Atlas
Preliminary Map
August 2009
From 861,488
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square