

Irania

Irania gutturalis

72 plots from 150 records

Virgin Yellow-vented Bulbul squares **18**

Single YvB month **37**

Two YvB months **34**

Seasonality

	presence	bs	eggs
January	38	-	-
February	28	-	-
March	32	-	-
April	3	-	-
May	-	-	-
June	-	-	-
July	-	-	-
August	-	-	-
September	-	-	-
October	-	-	-
November	1	-	-
December	48	-	-

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

VALUE

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

In a "normal" year this Palearctic migrant occurs rather sparsely throughout the drier thornbush habitat of the central plateau. In drier years, most probably when the short rains fail in eastern Kenya and NE Tanzania, greater numbers are found, especially in northern areas.

Despite the striking colours of adult males and the bright flash of orange underwings contrasting with that black tail it can be difficult to detect as it prefers quite dense shade.


The lack of records from the southern areas of the Masai Steppe is rather surprising and may reflect poor coverage.

with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
July 2006
From 722,000
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square



29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°

-1° -2° -3° -4° -5° -6° -7° -8° -9° 10° 11°