

# White-headed Black Chat

*Myrmecocichla arnoti*

412 plots from 734 records

The eastern and western populations of this Miombo endemic are clearly well separated by the Southern Highlands and the dry Bahi and Mtera Basins. The Miombo woodland along the Iringa-Dodoma escarpment and to the west and south of Bahi is now probably too dry for this species but may not have been in the recent past. The records from Mafia Is. is from 1937 and requires checking.

Virgin Yellow-vented Bulbul squares **12**

Single YvB month **23**

Two YvB months **22**

Seasonality  
presence bs eggs

Month	presence	bs	eggs
January	46	4	-
February	29	2	-
March	47	-	-
April	32	-	-
May	40	4	-
June	54	1	-
July	107	2	-
August	77	1	-
September	72	1	1
October	73	2	2
November	68	5	4
December	44	5	-
month 13	11		

## Status

- Presence
- Breeding season
- Month with eggs

## Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

Tanzania Bird Atlas  
Preliminary Map  
July 2009  
From 850,662  
Database records

J F M A  
M J J A  
S O N D  
# # # #

Monthly records  
within each square

