

# Karamoja Apalis

*Apalis karamojae*

24 plots from 30 records

Virgin Yellow-vented Bulbul squares **18**

Single YvB month **37**

Two YvB months **34**

Seasonality	presence		bs	eggs
January	2	-	-	-
February	2	-	-	-
March	-	-	-	-
April	-	-	-	-
May	2	-	-	-
June	4	-	-	-
July	8	-	-	-
August	4	-	1	-
September	1	-	-	-
October	1	-	-	-
November	2	-	-	-
December	3	-	-	-

The remarkable northward expansion of this species continues with recent records from the Mara GR in Kenya. It is now a regular sight in the western corridor of the Serengeti NP wherever mature Whistling Thorn occurs.

We need further records from the Wembere Steppe and adjacent drainage lines to fill that gap due west of lake Eyasi, one of the most poorly worked areas in the country.

That single record from the eastern border of Tarangire NP is a reminder of just how far this bird can move.

## Status

- Presence
- Breeding season
- Month with eggs

## Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

with support from  
THE LEVERHULME TRUST

**Tanzania Bird Atlas**  
**Preliminary Map**  
**May 2006**  
**From 722,040**  
**Database records**

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
within each square

