

# Upcher's Warbler

*Hippolais languida*

30 plots from 46 records

Virgin Yellow-vented Bulbul squares **13**

Single YvB month **25**

Two YvB months **28**

Seasonality  
presence bs eggs

Month	presence	bs	eggs
January	23	-	-
February	5	-	-
March	3	-	-
April	3	-	-
May	-	-	-
June	-	-	-
July	-	-	-
August	-	-	-
September	-	-	-
October	-	-	-
November	2	-	-
December	10	-	-

## Status

- Presence
- Breeding season
- Month with eggs

## Elevation (m)

### VALUE

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

One of the rarer Palearctic migrants at the southern end of its wintering range in NETanzania. The almost black tail contrasts well with a grey body and the tail wagging produces an elipse with the tip rather than the simpler up and down motion of a Olivaceous Warbler. Likely to be more common within this range than these few records suggest.

Tanzania Bird Atlas  
Preliminary Map  
May 2008  
From 807,507  
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
within each square

