

# Usambara Hyliota

## *Hyliota usambarae*

9 plots from 10 records

Virgin Yellow-vented Bulbul squares **14**

Single YvB month **30**

Two YvB months **31**

### Seasonality

	presence	bs	eggs
January	-	-	-
February	2	-	-
March	1	-	-
April	1	-	-
May	1	-	-
June	-	-	-
July	-	-	-
August	3	-	-
September	1	-	-
October	-	-	-
November	1	-	-
December	-	-	-

	presence	bs	eggs
January	-	-	-
February	2	-	-
March	1	-	-
April	1	-	-
May	1	-	-
June	-	-	-
July	-	-	-
August	3	-	-
September	1	-	-
October	-	-	-
November	1	-	-
December	-	-	-

### Status

- Presence
- Breeding season
- Month with eggs

### Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

This isolated population of the Southern Hyliota *H.australis* sometimes treated as a full species.


There is a single recent record from Msumbugwe FR south of the Pangani River and an old record from Ambangulu on the sw slopes of the West Usambara. All other records are from the East Usambara where it is an uncommon bird.

with support from  
THE LEVERHULME TRUST

**Tanzania Bird Atlas**  
**Preliminary Map**  
**April 2007**  
**From 754,349**  
**Database records**

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
within each square



29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°