

Yellow-bellied Hyliota

Hyliota flavigaster

152 plots from 221 records

Virgin Yellow-vented Bulbul squares **14**

Single YvB month **30**

Two YvB months **31**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	5	-	-
February	12	-	-
March	20	-	-
April	16	-	-
May	17	-	-
June	7	-	-
July	36	-	-
August	24	-	-
September	37	-	-
October	18	-	-
November	18	2	-
December	9	-	-

Status


- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

A widespread low density species throughout the Miombo belt. Most bird parties have a pair of these birds in them but as they are rather quiet and forage in the canopy they are easily missed.

The record from 3101B is from Minziro forest and may be of the race *slatini* of *H. australis* the Southern Hyliota.



with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
April 2007
From 754,349
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°