

Western Yellowbill

Ceuthmochares aereus

41 plots plots from 77 records

Virgin Yellow-vented Bulbul squares **11**

Single YvB month **21**

Two YvB months **25**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	6	-	-
February	1	-	-
March	8	-	-
April	7	-	-
May	-	-	-
June	4	-	-
July	5	-	-
August	9	-	-
September	7	-	-
October	8	-	-
November	11	1	1
December	4	-	-
month 13	5	-	-

Status


- Presence
- Breeding season
- Month with eggs

30m SRTM

- 0 - 100
- 100 - 200
- 200 - 300
- 300 - 400
- 400 - 450
- 450 - 500
- 500 - 550
- 550 - 600
- 600 - 650
- 650 - 700
- 700 - 750
- 750 - 800
- 800 - 850
- 850 - 900
- 900 - 950
- 950 - 1,000
- 1,000 - 1,050
- 1,050 - 1,100
- 1,100 - 1,200
- 1,200 - 1,300
- 1,300 - 1,400
- 1,400 - 1,500
- 1,500 - 1,600
- 1,600 - 1,700
- 1,700 - 1,800
- 1,800 - 1,900
- 1,900 - 2,000
- 2,000 - 2,100
- 2,100 - 2,200
- 2,200 - 2,300
- 2,300 - 2,400
- 2,400 - 2,500

A widespread West and central African bird at the eastern edge of its extensive range. The paucity of data reflects the scarcity of observers in extreme western Tanzania.

The range may well be continuous as extensive riverine habitat still exists throughout this area despite the severe local degradation caused by waves of refugees from DRC, Burundi and Rwanda.



Tanzania Bird Atlas
Preliminary Map
October 2009
From 868,376
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square