

Common Whitethroat

Sylvia communis

173 plots from 729 records

A locally common migrant and winter visitor in the dry central plateau and southern highland areas but rare to absent in the western and eastern Miombo woodlands. The paucity of records in the southern Masai Steppe could be the result of poor coverage.

Virgin Yellow-vented Bulbul squares **33**

Single YvB month **34**

Two YvB months **30**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	74	-	-
February	50	-	-
March	282	-	-
April	143	-	-
May	6	-	-
June	-	-	-
July	-	-	-
August	-	-	-
September	-	-	-
October	1	-	-
November	47	-	-
December	126	-	-

Protected Areas

- National Park
- Game Reserve
- Conservaiona Area

Elevation (m)


- 1 - 300
- 300 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 2,000
- 2,000 - 2,300
- 2,300 - 2,600

with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
December 2004
From 649,600
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square



29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°

-1°
-2°
-3°
-4°
-5°
-6°
-7°
-8°
-9°
-10°
-11°