

Pangani Longclaw

Macronyx aurantiigula

175 plots from 470 records

Virgin Yellow-vented Bulbul squares **14**

Single YvB month **30**

Two YvB months **31**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	48	2	2
February	45	2	1
March	39	2	-
April	33	5	1
May	19	2	2
June	36	1	-
July	45	-	-
August	37	1	-
September	28	-	-
October	32	1	-
November	39	3	-
December	43	2	-

Status

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

The westwards expansion over the Rift valley wall is quite recent, the first record was by John Beesley in 1981. The paucity of records on the Masai Steppe is puzzling and it surely occurs more often in eastern Mkomzai than suggested here. There does not appear to be any records below 300m.

It should be looked for in the Eyasi Basin especially on the Wembere Steppe and in the vicinity of Lake Kitangire. It would be interesting to monitor a continued expansion westwards and to the SW.

The red squares show range overlap (at the square level) with Yellow-throated Longclaw.

with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
May 2007
From 762,609
Database records

J F M A
M J J A
S O N D
#

Monthly records
within each square

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°