

Ashy Starling

Cosmopsarus unicolor

481 plots from 1,776 records

Virgin Yellow-vented Bulbul squares **16**

Single YvB month **33**

Two YvB months **33**

Seasonality

	presence	bs	eggs
January	134	6	-
February	121	4	1
March	149	6	2
April	119	9	2
May	58	2	-
June	138	-	-
July	189	1	-
August	213	-	-
September	128	-	-
October	226	-	-
November	116	-	-
December	152	-	-

Species

- Presence
- Breeding season
- Month with eggs

Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- 1,900 - 2,100

White-winged Starling shown by red crosses

The White-winged Starling is endemic to Miombo Woodland with disjunct populations in Tanzania, Zambia and Angola. Ashy Starlings avoid Miombo Woodland but overlap with WwS at the level of the Atlas square where Acacia woodland occurs along drainage lines within the Miombo belt.

Note that this only occurs where these drainage lines flow into river systems within the Acacia-Comiphora zones of the central plateau and not into the isolated drainage basins of Lake Rukwa and the Malagarasi system.

The western limits of Ashy Starling will be defined by contiguous belts of Miombo, by soil types and watersheds rather than altitude and rainfall which help define the eastern, southern and northern limits.

with support from
THE LEVERHULME TRUST

Tanzania Bird Atlas
Preliminary Map
October 2006
From 730,163
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

